

3d803 U.S. PTC

UTILITY
PATENT APPLICATION
TRANSMITTAL

for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.

2314-206

Total Pages

First Named Inventor or Application Identifier

Ann H. CORNELL-BELL

Express Mail Label No.

914 U.S. PTO
09/666837
09/21/80

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner of Patents
Box Patent Application
Washington, D.C. 20231

1. ☐ ☐ Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)
2. ☐ ☒ Specification Total pages [31]
(preferred arrangement set forth below)
- Descriptive title of the invention
 - Cross references to Related Applications
 - Statement Regarding Fed sponsored R&D
 - Reference to Microfiche Appendix
 - Background of the Invention
 - Brief Summary of the Invention
 - Brief Description of the Drawings
 - Detailed Description
 - Claims
 - Abstract of the Disclosure
3. ☐ ☒ Drawing(s) (35 USC 113) (Total Sheets) [6]
4. ☐ ☐ Oath or Declaration (Total Pages) []
- a. ☐ ☐ Newly executed (original or copy)
 - b. ☐ ☐ Copy from a prior application
(37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
- [Note Box 5 below]**
- i. ☐ ☐ **DELETION OF INVENTOR(S)**
Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b)
5. ☐ ☐ Incorporation by Reference (useable if Box 4b is checked) The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. [] Microfiche Computer Program (Appendix)
7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - a. [XX] Computer Readable Copy
 - b. [XX] Paper Copy (identical to computer copy) (8 pages)
 - c. [XX] Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. [] Assignment Papers (cover sheet & documents)
9. [] 37 CFR 3.73(b) Statement
(when there is an assignee)
[] Power of Attorney
10. [] English Translation Document (if applicable)
11. [] Information Disclosure Statement /PTO 1449
[] Copies of IDS Citations
12. [] Preliminary Amendment
13. [X] Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
14. [] Small Entity Statement(s)
[] Statement Filed in prior application,
Status still proper and desired
15. [] Certified Copy of Priority Document(s).
(if foreign priority is claimed)
16. [] Other:

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.:

18. CORRESPONDENCE ADDRESS

[1 Customer Number or Bar Code Label

or [X] Correspondence address below

(Insert Customer No. or Attach bar code label here)

Name _____

Jeffrey L. Ihnen, Reg. No. 28,957

Address

Rothwell, Figg, Ernst & Manbeck
Suite 701-East, 555 13th Street, N.W.

City

Washington

State

D.C.

Zip Code

20004

Country

U.S.A.

Telephone

202-783-6040

Fax

202-783-6031

TITLE OF THE INVENTION

USES OF KAPPA-CONOTOXIN PVIIA

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is related to U.S. provisional patent application Serial No. 60/219,438 filed on 20 July 2000 and to U.S. provisional patent application Serial No. 60/155,135 filed on 22 September 1999, each of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

The invention relates to uses of kappa-conotoxin PVIIA (κ -PVIIA), analogs and derivatives for activating (i.e., opening) ATP-sensitive K^+ channels. The activation of ATP-sensitive K^+ channels is useful for treating many physiological disorders, as described in further detail herein.

The publications and other materials used herein to illuminate the background of the invention, and in particular, cases to provide additional details respecting the practice, are incorporated by reference, and for convenience are referenced in the following text by author and date and are listed alphabetically by author in the appended bibliography.

κ -PVIIA, a 27 amino acid peptide that was originally purified from the venom of the purple cone snail *Conus purpurascens* (Terlau et al., 1996; US Patent No. 5,672,682) has been previously identified as a potent antagonist of the *Shaker* H4 potassium channel ($IC_{50} \sim 60$ nM). In the same study, no detectable activity on the voltage-gated potassium channels Kv1.1 or Kv1.4 (Terlau et al., 1996) was noted. Chimeras constructed from the *Shaker* and the Kv1.1 K^+ channels have identified the putative pore-forming region between the fifth and sixth transmembrane region as the site of the toxin sensitivity (Shon et al., 1998). It appears that κ -PVIIA interacts with the external tetraethylammonium binding site on the *Shaker* channel. Although both κ -PVIIA and charybdotoxin inhibit the *Shaker* channel, they must interact differently. The F425G *Shaker* mutation increases charybdotoxin affinity by three orders of magnitude but abolishes κ -PVIIA sensitivity (Shon et al., 1998). κ -PVIIA appears to block the ion pore with a 1:1 stoichiometry, and its binding to open or closed channels is very different (Terlau et al., 1999). Chronically applied to whole oocytes or outside-out patches, kappa-PVIIA inhibition appears as a voltage-dependent relaxation in response to the depolarizing pulse used to activate the channels (Garcia et al., 1999).

Potassium channels are vital in controlling the resting membrane potential in excitable cells and can be broadly subdivided into three classes, voltage-gated channels, Ca^{2+} activated channels and ATP-sensitive K^+ channels. ATP-sensitive potassium channels were originally described in

cardiac tissue (Noma, 1983). In subsequent years they have also been identified in pancreatic cells, skeletal, vascular and neuronal tissue. This group of K^+ channels are modulated by intracellular ATP levels and as such, couple cellular metabolism to electrical activity. Enhanced levels of ATP produce closure of the K_{ATP} channels. The K_{ATP} channel is thought to be an octomeric complex comprised of two different subunits in a 1:1 stoichiometry; a weakly inward rectifying K^+ channel Kir6.x (6.1 or 6.2), which is thought to form the channel pore, and a sulphonylurea (SUR) subunit. So far, three variants of the SUR have been identified: SUR1, SUR2A and SUR2B. While the Kir6.2 subunit is common to K_{ATP} channels in cardiac, pancreatic and neuronal tissue (Kir6.1 is preferentially expressed in vascular smooth muscle tissue), the SUR is differentially expressed. Kir6.2/SUR1 reconstitute the neuronal/pancreatic beta-cell channel, whereas Kir6.2/SUR2A are proposed to reconstitute the cardiac K_{ATP} channels.

Potassium channels comprise a large and diverse group of proteins that, through maintenance of the cellular membrane potential, are fundamental in normal biological function. The potential therapeutic applications for compounds that open K^+ channels are far-reaching and include treatments of a wide range of disease and injury states, including cerebral and cardiac ischemia and asthma. Recently, considerable interest has focused around the ability of K^+ channel openers to produce relaxation of airway smooth muscle, and as such, these compounds may offer a novel approach to the treatment of bronchial asthma (Lin et al., 1998; Muller-Schweinitzer and Fozard, 1997; Morley, 1994; Barnes, 1992). Furthermore, the cardioprotective effects of K^+ channel openers are now well established in experimental animal models of cardiac ischemia (Jung et al., 1998; Kouchi et al., 1998). Less is known about the ability of these compounds to limit neuronal damage caused from cerebral ischemia. Most progress in the treatment of cerebral ischemia has focused around the development of compounds to reduce the influx of sodium and calcium ions. K^+ channel openers, which restore the resting membrane potential, could also be employed to reduce acute damage associated with an ischemic episode in neuronal tissue (Reshef et al., 1998; Wind et al., 1997), as well as reducing glutamate-induced excitotoxicity (Lauritzen et al., 1997). However, clinical use of K_{ATP} openers has been somewhat limited due to their cardiovascular side effects (i.e., drop in blood pressure).

Thus, it is desired to develop new agents for opening K_{ATP} channels which can be used to treat a wide range of disease and injury states, including cerebral and cardiac ischemia and asthma.

SUMMARY OF THE INVENTION

The invention relates to uses of kappa-conotoxin PVIIA (κ -PVIIA), analogs and derivatives for activating ATP-sensitive K^+ channels. The opening of ATP-sensitive K^+ channels is useful for treating many physiological disorders as described in further detail herein.

More specifically, the present invention is directed to the use of κ -PVIIA, its analogs, derivatives and physiologically acceptable salts thereof for opening K_{ATP} channels which can be used to treat cardiac ischemia, neuronal ischemia, ocular ischemia and asthma.

For purposes of the present invention, κ -PVIIA refers to a peptide having the following general formula:

Cys-Xaa₁-Ile-Xaa₂-Asn-Gln-Xaa₃-Cys-Xaa₄-Gln-Xaa₅-Leu-Asp-Asp-Cys-Cys-Ser-Xaa₁-Xaa₃-Cys-Asn-Xaa₁-Xaa₄-Asn-Xaa₃-Cys-Val (SEQ ID NO:1), wherein Xaa₁ and Xaa₃ are independently Arg, homoarginine, ornithine, Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys, any synthetic basic amino acid, His or halo-His; Xaa₂ is Pro or hydroxy-Pro (Hyp); Xaa₄ is Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; and Xaa₅ is His or halo-His. The C-terminus may contain a free carboxyl group or an amide group. The halo is preferably bromine, chlorine or iodine. It is preferred that Xaa₁ is Arg and Xaa₅ is His. It is more preferred that Xaa₁ is Arg, Xaa₃ is Lys, Xaa₄ is Phe and Xaa₅ is His. It is further preferred that the C-terminus contains a free carboxyl group.

The κ -PVIIA analogs refer to peptides having the following formulas:

κ -PVIIA[R18A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Ala-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:2);

κ -PVIIA[R22A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Ala-Phe-Asn-Lys-Cys-Val (SEQ ID NO:3);

κ -PVIIA[I3A]: Cys-Arg-Ala-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:4);

κ -PVIIA[K19A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Ala-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:5);

κ -PVIIA[R2A]: Cys-Ala-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:6);

κ -PVIIA[F9A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Ala-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:7);

κ-PVIA[K25A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Ala-Cys-Val (SEQ ID NO:8);

κ-PVIA[R2K]: Cys-Lys-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:9);

κ-PVIA[K7A]: Cys-Arg-Ile-Hyp-Asn-Gln-Ala-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:10);

κ-PVIA[F9M]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Met-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:11);

κ-PVIA[F9Y]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Tyr-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:12);

κ-PVIA[R2Q]: Cys-Gln-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:13);

κ-PVIA[H11A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-Ala-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:14);

κ-PVIA[D14A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Ala-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:15);

κ-PVIA[Q6A]: Cys-Arg-Ile-Hyp-Asn-Ala-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:16);

κ-PVIA[N21A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Ala-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:17);

κ-PVIA[S17A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ala-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:18);

κ-PVIA[N24A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Ala-Lys-Cys-Val (SEQ ID NO:19);

κ-PVIA[L12A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Ala-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:20);

κ-PVIA[D13A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Ala-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:21);

κ-PVIA[Q10A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Ala-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:22);

κ-PVIA[V27A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Ala (SEQ ID NO:23);

κ-PVIIA[O4A]: Cys-Arg-Ile-Ala-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:24); and

κ-PVIIA[N5A]: Cys-Arg-Ile-Hyp-Ala-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-
Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:25).

It is preferred that the C-terminus contains a free carboxyl group.

The present invention further relates to derivatives of the above peptides in which the Arg residues may be substituted by Lys, ornithine, homoarginine, nor-Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys or any synthetic basic amino acid; the Lys residues may be substituted by Arg, ornithine, homoarginine, nor-Lys, or any synthetic basic amino acid; the Tyr residues may be substituted with any synthetic hydroxy containing amino acid; the Ser residues may be substituted with Thr or any synthetic hydroxylated amino acid; the Thr residues may be substituted with Ser or any synthetic hydroxylated amino acid; the Phe and Trp residues may be substituted with any synthetic aromatic amino acid; and the Asn, Ser, Thr or Hyp residues may be glycosylated (contain an N-glycan or an O-glycan). The Cys residues may be in D or L configuration and may optionally be substituted with homocysteine (D or L). The Tyr residues may also be substituted with the 3-hydroxyl or 2-hydroxyl isomers (meta-Tyr or ortho-Tyr, respectively) and corresponding O-sulpho- and O-phospho-derivatives. The acidic amino acid residues may be substituted with any synthetic acidic amino acid, e.g., tetrazolyl derivatives of Gly and Ala. The aliphatic amino acids may be substituted by synthetic derivatives bearing non-natural aliphatic branched or linear side chains C_nH_{2n+2} up to and including $n=8$.

Examples of synthetic aromatic amino acid include, but are not limited to, such as nitro-Phe, 4-substituted-Phe wherein the substituent is C_1 - C_3 alkyl, carboxyl, hydroxymethyl, sulphomethyl, halo, phenyl, -CHO, -CN, -SO₃H and -NHAc. Examples of synthetic hydroxy containing amino acid, include, but are not limited to, such as 4-hydroxymethyl-Phe, 4-hydroxyphenyl-Gly, 2,6-dimethyl-Tyr and 5-amino-Tyr. Examples of synthetic basic amino acids include, but are not limited to, N-1-(2-pyrazolyl)-Arg, 2-(4-piperinyl)-Gly, 2-(4-piperinyl)-Ala, 2-[3-(2S)pyrrolinyl]-Gly and 2-[3-(2S)pyrrolinyl]-Ala. These and other synthetic basic amino acids, synthetic hydroxy containing amino acids or synthetic aromatic amino acids are described in Building Block Index, Version 3.0 (1999 Catalog, pages 4-47 for hydroxy containing amino acids and aromatic amino acids and pages 66-87 for basic amino acids; see also <http://www.amino-acids.com>), incorporated herein by reference, by and available from RSP Amino Acid Analogues, Inc., Worcester, MA. The residues containing protecting groups are deprotected using conventional techniques. Examples of

synthetic acid amino acids include those derivatives bearing acidic functionality, including carboxyl, phosphate, sulfonate and synthetic tetrazolyl derivatives such as described by Ornstein et al. (1993) and in U.S. Patent No. 5,331,001, each incorporated herein by reference.

In accordance with the present invention, a glycan shall mean any N-, S- or O-linked mono-, di-, tri-, poly- or oligosaccharide that can be attached to any hydroxy, amino or thiol group of natural or modified amino acids by synthetic or enzymatic methodologies known in the art. The monosaccharides making up the glycan can include D-allose, D-altrose, D-glucose, D-mannose, D-gulose, D-idose, D-galactose, D-talose, D-galactosamine, D-glucosamine, D-N-acetyl-glucosamine (GlcNAc), D-N-acetyl-galactosamine (GalNAc), D-fucose or D-arabinose. These saccharides may be structurally modified, e.g., with one or more O-sulfate, O-phosphate, O-acetyl or acidic groups, such as sialic acid, including combinations thereof. The glycan may also include similar polyhydroxy groups, such as D-penicillamine 2,5 and halogenated derivatives thereof or polypropylene glycol derivatives. The glycosidic linkage is beta and 1-4 or 1-3, preferably 1-3. The linkage between the glycan and the amino acid may be alpha or beta, preferably alpha and is 1-.

Core O-glycans have been described by Van de Steen et al. (1998), incorporated herein by reference. Mucin type O-linked oligosaccharides are attached to Ser or Thr (or other hydroxylated residues of the present peptides) by a GalNAc residue. The monosaccharide building blocks and the linkage attached to this first GalNAc residue define the "core glycans," of which eight have been identified. The type of glycosidic linkage (orientation and connectivities) are defined for each core glycan. Suitable glycans and glycan analogs are described further in U.S. Serial No. 09/420,797, filed 19 October 1999 and in PCT Application No. PCT/US99/ 24380, filed 19 October 1999 (PCT Published Application No. WO 00/23092), each incorporated herein by reference. A preferred glycan is Gal(β 1-3)GalNAc(α 1-).

Optionally, in the above peptides, pairs of Cys residues may be replaced pairwise with isoteric lactam or ester-thioether replacements, such as Ser/(Glu or Asp), Lys/(Glu or Asp) or Cys/Ala combinations. Sequential coupling by known methods (Barnay et al., 2000; Hruby et al., 1994; Bitan et al., 1997) allows replacement of native Cys bridges with lactam bridges. Thioether analogs may be readily synthesized using halo-Ala residues commercially available from RSP Amino Acid Analogues.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows fluorimetry measurements of intracellular K^+ (determined with PBFI dye) following exposure to increasing concentrations of κ -PVIIA in primary cultures of ventricular myocytes. The data shown is from one trial and is represented as mean change in fluorescence \pm S.E.M. (* $p < 0.05$, unpaired t-test).

Figures 2A-2B show fluorimetry measurements of membrane potential (determined with Di-8-ANEPPs dye) following exposure to increasing concentrations of κ -PVIIA in primary cultures of ventricular myocytes (Fig. 2A) or cortex (Fig. 2B). Cells were loaded into 96 well plates at least six days before the experiment. Results are expressed as Mean \pm SEM and represent average data from between two and five individual trials.

Figures 3A-3B are bar graphs showing the inhibition of the κ -PVIIA (100nM) response with 10nM Glibenclamide (Glib) in primary cultures of myocytes (Fig. 3A) or with 50uM Tolbutamide (Tolb) in primary cultures of cortex (Fig. 3B). Data represents mean \pm S.E.M.

Figures 4A-4C are whole cell recordings showing currents elicited by κ -PVIIA in (Fig. 4A) cortical cells and (Fig. 4B) myocytes. Fig 4C shows I-V relationship of κ -PVIIA-induced current from a cardiac myocyte.

Figure 5 is a bar graph showing the protective effect of 10 nM κ -PVIIA against hypoxia induced depolarization. Bars represent Mean \pm S.E.M.

Figure 6 shows the effect of increasing concentrations of κ -PVIIA on glutamate-induced (100uM) excitotoxicity measured six hours following glutamate washout (three to six trials).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The invention relates to uses of kappa-conotoxin PVIIA (κ -PVIIA), analogs and derivatives for activating ATP-sensitive K^+ channels. The activation of ATP-sensitive K^+ channels is useful for treating many physiological disorders, as described in further detail herein.

More specifically, the present invention is directed to the use of κ -PVIIA, analogs and derivatives for opening K_{ATP} channels which can be used to treat cardiac ischemia, neuronal ischemia, ocular ischemia and asthma.

The present invention, in another aspect, relates to a pharmaceutical composition comprising an effective amount of κ -PVIIA, analogs, derivatives or pharmaceutically acceptable salts. Such a pharmaceutical composition has the capability of acting as an activator for K_{ATP} channels. Thus,

the pharmaceutical compositions of the present invention are useful in the treatment of the disorders noted above.

κ-PVIIA can be isolated from *Conus purpurascens* as described in U.S. Patent No. 5,672,682, or it can be chemically synthesized by general synthetic methods such as described in U.S. Patent No. 5,672,682. Alternatively, the native peptide can be synthesized by conventional recombinant DNA techniques (Sambrook et al., 1989) using the DNA encoding κ-PVIIA (Shon et al., 1998). The peptides are also synthesized using an automated synthesizer. Amino acids are sequentially coupled to an MBHA Rink resin (typically 100 mg of resin) beginning at the C-terminus using an Advanced ChemTech 357 Automatic Peptide Synthesizer. Couplings are carried out using 1,3-diisopropylcarbodiimide in N-methylpyrrolidinone (NMP) or by 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and diethylisopropylethylamine (DIEA). The Fmoc protecting group is removed by treatment with a 20% solution of piperidine in dimethylformamide (DMF). Resins are subsequently washed with DMF (twice), followed by methanol and NMP.

Mutens, analogs or active fragments, of the foregoing τ-conotoxin peptides are also contemplated here. See, e.g., Hammerland et al (1992). Derivative mutens, analogs or active fragments of the conotoxin peptides may be synthesized according to known techniques, including conservative amino acid substitutions, such as outlined in U.S. Patents No. 5,545,723 (see particularly col. 2, line 50 to col. 3, line 8); 5,534,615 (see particularly col. 19, line 45 to col. 22, line 33); and 5,364,769 (see particularly col. 4, line 55 to col. 7, line 26), each incorporated herein by reference.

Pharmaceutical compositions containing a compound of the present invention or its pharmaceutically acceptable salts as the active ingredient can be prepared according to conventional pharmaceutical compounding techniques. See, for example, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990, Mack Publishing Co., Easton, PA). Typically, a K_{ATP} channel activating amount of the active ingredient will be admixed with a pharmaceutically acceptable carrier. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., intravenous, oral or parenteral. The compositions may further contain antioxidantizing agents, stabilizing agents, preservatives and the like. For examples of delivery methods, see U.S. Patent No. 5,844,077, incorporated herein by reference.

"Pharmaceutical composition" means physically discrete coherent portions suitable for medical administration. "Pharmaceutical composition in dosage unit form" means physically

discrete coherent units suitable for medical administration, each containing a daily dose or a multiple (up to four times) or a sub-multiple (down to a fortieth) of a daily dose of the active compound in association with a carrier and/or enclosed within an envelope. Whether the composition contains a daily dose, or for example, a half, a third or a quarter of a daily dose, will depend on whether the pharmaceutical composition is to be administered once or, for example, twice, three times or four times a day, respectively.

The term "salt", as used herein, denotes acidic and/or basic salts, formed with inorganic or organic acids and/or bases, preferably basic salts. While pharmaceutically acceptable salts are preferred, particularly when employing the compounds of the invention as medicaments, other salts find utility, for example, in processing these compounds, or where non-medicament-type uses are contemplated. Salts of these compounds may be prepared by art-recognized techniques.

Examples of such pharmaceutically acceptable salts include, but are not limited to, inorganic and organic addition salts, such as hydrochloride, sulphates, nitrates or phosphates and acetates, trifluoroacetates, propionates, succinates, benzoates, citrates, tartrates, fumarates, maleates, methane-sulfonates, isothionates, theophylline acetates, salicylates, respectively, or the like. Lower alkyl quaternary ammonium salts and the like are suitable, as well.

As used herein, the term "pharmaceutically acceptable" carrier means a non-toxic, inert solid, semi-solid liquid filler, diluent, encapsulating material, formulation auxiliary of any type, or simply a sterile aqueous medium, such as saline. Some examples of the materials that can serve as pharmaceutically acceptable carriers are sugars, such as lactose, glucose and sucrose, starches such as corn starch and potato starch, cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt, gelatin, talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol, polyols such as glycerin, sorbitol, mannitol and polyethylene glycol; esters such as ethyl oleate and ethyl laurate, agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline, Ringer's solution; ethyl alcohol and phosphate buffer solutions, as well as other non-toxic compatible substances used in pharmaceutical formulations.

Wetting agents, emulsifiers and lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator. Examples of pharmaceutically acceptable antioxidants include,

but are not limited to, water soluble antioxidants such as ascorbic acid, cysteine hydrochloride, sodium bisulfite, sodium metabisulfite, sodium sulfite, and the like; oil soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol and the like; and the metal chelating agents such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid and the like.

For oral administration, the compounds can be formulated into solid or liquid preparations such as capsules, pills, tablets, lozenges, melts, powders, suspensions or emulsions. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, suspending agents and the like in the case of oral liquid preparations (such as, for example, suspensions, elixirs and solutions); or carriers such as starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations (such as, for example, powders, capsules and tablets). Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar-coated or enteric-coated by standard techniques. The active agent can be encapsulated to make it stable for passage through the gastrointestinal tract, while at the same time allowing for passage across the blood brain barrier. See for example, WO 96/11698.

For parenteral administration, the compound may be dissolved in a pharmaceutical carrier and administered as either a solution or a suspension. Illustrative of suitable carriers are water, saline, dextrose solutions, fructose solutions, ethanol, or oils of animal, vegetative or synthetic origin. The carrier may also contain other ingredients, for example, preservatives, suspending agents, solubilizing agents, buffers and the like. When the compounds are being administered intrathecally, they may also be dissolved in cerebrospinal fluid.

A variety of administration routes are available. The particular mode selected will depend of course, upon the particular drug selected, the severity of the disease state being treated and the dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, sublingual, topical, nasal, transdermal or parenteral routes. The term "parenteral" includes subcutaneous, intravenous, epidural, irrigation, intramuscular, release pumps, or infusion.

For example, administration of the active agent according to this invention may be achieved using any suitable delivery means, including:

- (a) pump (see, e.g., Lauer & Hatton (1993), Zimm et al. (1984) and Ettinger et al. (1978));
- (b), microencapsulation (see, e.g., U.S. Patent Nos. 4,352,883; 4,353,888; and 5,084,350);
- (c) continuous release polymer implants (see, e.g., U.S. Patent No. 4,883,666);
- (d) macroencapsulation (see, e.g., U.S. Patent Nos. 5,284,761, 5,158,881, 4,976,859 and 4,968,733 and published PCT patent applications WO92/19195, WO 95/05452);
- (e) naked or unencapsulated cell grafts to the CNS (see, e.g., U.S. Patent Nos. 5,082,670 and 5,618,531);
- (f) injection, either subcutaneously, intravenously, intra-arterially, intramuscularly, or to other suitable site; or

(g) oral administration, in capsule, liquid, tablet, pill, or prolonged release formulation.

In one embodiment of this invention, an active agent is delivered directly into the CNS, preferably to the brain ventricles, brain parenchyma, the intrathecal space or other suitable CNS location, most preferably intrathecally.

Alternatively, targeting therapies may be used to deliver the active agent more specifically to certain types of cells, by the use of targeting systems such as antibodies or cell-specific ligands. Targeting may be desirable for a variety of reasons, e.g. if the agent is unacceptably toxic, if it would otherwise require too high a dosage, or if it would not otherwise be able to enter target cells.

The active agents, which are peptides, can also be administered in a cell based delivery system in which a DNA sequence encoding an active agent is introduced into cells designed for implantation in the body of the patient, especially in the spinal cord region. Suitable delivery systems are described in U.S. Patent No. 5,550,050 and published PCT Application Nos. WO 92/19195, WO 94/25503, WO 95/01203, WO 95/05452, WO 96/02286, WO 96/02646, WO 96/40871, WO 96/40959 and WO 97/12635. Suitable DNA sequences can be prepared synthetically for each active agent on the basis of the developed sequences and the known genetic code.

The active agent is preferably administered in an therapeutically effective amount. By a "therapeutically effective amount" or simply "effective amount" of an active compound is meant a sufficient amount of the compound to treat or alleviate pain or to induce analgesia at a reasonable benefit/risk ratio applicable to any medical treatment. The actual amount administered, and the rate and time-course of administration, will depend on the nature and severity of the condition being treated. Prescription of treatment, e.g. decisions on dosage, timing, etc., is within the responsibility

of general practitioners or specialists, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of techniques and protocols can be found in *Remington's Pharmaceutical Sciences*.

Dosage may be adjusted appropriately to achieve desired drug levels, locally or systemically. Typically, the active agents of the present invention exhibit their effect at a dosage range of from about 0.001 mg/kg to about 250 mg/kg, preferably from about 0.01 mg/kg to about 100 mg/kg, of the active ingredient and more preferably, from about 0.05 mg/kg to about 75 mg/kg. A suitable dose can be administered in multiple sub-doses per day. Typically, a dose or sub-dose may contain from about 0.1 mg to about 500 mg of the active ingredient per unit dosage form. A more preferred dosage will contain from about 0.5 mg to about 100 mg of active ingredient per unit dosage form. Dosages are generally initiated at lower levels and increased until desired effects are achieved.

Advantageously, the compositions are formulated as dosage units, each unit being adapted to supply a fixed dose of active ingredients. Tablets, coated tablets, capsules, ampoules and suppositories are examples of dosage forms according to the invention.

It is only necessary that the active ingredient constitute an effective amount, i.e., such that a suitable effective dosage will be consistent with the dosage form employed in single or multiple unit doses. The exact individual dosages, as well as daily dosages, are determined according to standard medical principles under the direction of a physician or veterinarian for use humans or animals.

The pharmaceutical compositions will generally contain from about 0.0001 to 99 wt. %, preferably about 0.001 to 50 wt. %, more preferably about 0.01 to 10 wt.% of the active ingredient by weight of the total composition. In addition to the active agent, the pharmaceutical compositions and medicaments can also contain other pharmaceutically active compounds. Examples of other pharmaceutically active compounds include, but are not limited to, analgesic agents, cytokines and therapeutic agents in all of the major areas of clinical medicine. When used with other pharmaceutically active compounds, the conotoxin peptides of the present invention may be delivered in the form of drug cocktails. A cocktail is a mixture of any one of the compounds useful with this invention with another drug or agent. In this embodiment, a common administration vehicle (e.g., pill, tablet, implant, pump, injectable solution, etc.) would contain both the instant composition in combination supplementary potentiating agent. The individual drugs of the cocktail are each administered in therapeutically effective amounts. A therapeutically effective amount will

be determined by the parameters described above; but, in any event, is that amount which establishes a level of the drugs in the area of body where the drugs are required for a period of time which is effective in attaining the desired effects.

Activators of K_{ATP} channels have therapeutic significance for the treatment of asthma, cardiac ischemia and cerebral ischemia, among others.

Asthma: Asthma is a serious and common condition that effects approximately 12 million people in the United States alone. This disorder is particularly serious in children and it has been estimated that the greatest number of asthma patients are those under the age of 18 (National Health Survey, National Center of Health Statistics, 1989). The disease is characterized by chronic inflammation and hyper-responsiveness of the airway which results in periodic attacks of wheezing and difficulty in breathing. An attack occurs when the airway smooth muscle become inflamed and swells as a result of exposure to a trigger substance. In severe cases, the airway may become blocked or obstructed as a result of the smooth muscle contraction. Further exacerbating the problem is the release of large quantities of mucus which also act to block the airway. Chronic asthmatics are most commonly treated prophylactically with inhaled corticosteroids and acutely with inhaled bronchodilators, usually β -2 agonists. However, chronic treatment with inhaled corticosteroids has an associated risk of immune system impairment, hypertension, osteoporosis, adrenal gland malfunction and an increased susceptibility to fungal infections (Rakel, 1997). In addition use of β -2 agonists has been reported in some cases to cause adverse reactions including tremor, tachycardia and palpitations and muscle cramps (Rakel, 1997). Therefore, there is great potential in developing anti-asthmatic agents with fewer side-effects.

K^+ channel openers have been shown to be effective relaxants of airway smooth muscle reducing hyperactivity induced obstruction of intact airway. In cryopreserved human bronchi (Muller-Schweinitzer and Fozard, 1997) and in the isolated guinea pig tracheal preparation (Lin et al, 1998; Ando et al., 1997; Nielson-Kudsk, 1996; Nagai et al., 1991). K_{ATP} openers produced relaxation whether the muscle was contracted spontaneously or induced by a range of spasmogens. Under these conditions, the K^+ channel openers are thought to be acting to produce a K^+ ion efflux and consequent membrane hyperpolarization. As a result, voltage-sensitive Ca^{2+} channels would close and intracellular calcium levels would drop, producing muscular relaxation. The development of new and more specific K_{ATP} openers may offer a novel approach both to the prophylactic and symptomatic treatment of asthma.

K_{ATP} channels are present in many tissue types beyond just the target tissue, therefore their activation may result in unwanted side effects. In particular, as K_{ATP} channels are found in vascular smooth muscle, it is possible that in addition to the beneficial anti-asthmatic properties of K_{ATP} openers there could be an associated drop in blood pressure. It is possible that delivering the compound in inhalant form directly to the airway smooth muscle will allow the concentration of the compound to be reduced significantly thereby minimizing adverse reactions.

Cardiac Ischemia: While numerous subtypes of potassium channels in cardiac tissue have not yet been fully characterized, openers of K_{ATP} channels show great promise as cardioprotective agents. The beneficial vasodilatory effects afforded by K^+ channel openers in patients with angina pectoris are now well established (Chen et al., 1997; Goldschmidt et al., 1996; Yamabe et al., 1995; Koike et al., 1995). Furthermore, the activation of K_{ATP} channels appears also to be involved in the acute preconditioning of the myocardium following brief ischemic periods, acting to reduce the risk (Pell et al., 1998) and size of the reperfusion infarct (Kouchi et al., 1998).

Direct evidence for the cytoprotective properties of K_{ATP} channels was demonstrated by Jovanovic et al. (1998a). In these studies, the DNA encoding for the Kir6.2/SUR2A (cardiac K_{ATP}) channel were transfected in COS-7 monkey cells and the degree of calcium loading monitored. Untransfected cells were demonstrated to be vulnerable to the increases in intracellular calcium seen following hypoxia/reoxygenation. However, the transfection of the cells with the K_{ATP} channel conferred resistance to the potentially damaging effects of the hypoxia-reoxygenation. Thus, the cardiac K_{ATP} channels are likely to play a significant role in protecting the myocardium against reperfusion injury.

Cerebral Ischemia: Although treatment of cerebral ischemia has advanced significantly over the past 30 years, cerebral ischemia (stroke) still remains the third leading cause of death in the United States. More than 500,000 new stroke/ischemia cases are reported each year. Even though initial mortality is high (38%), there are close to three million survivors of stroke in the United States, and yearly cost for rehabilitation of these patients in the United States is close to \$17 billion (Rakel, 1997).

The initial cellular effects occur very rapidly (a matter of minutes) after an ischemic episode, whereas the actual cellular destruction does not occur until several hours or days following the infarction. Initial effects include depolarization due to bioenergetic failure, and inactivation of Na^+ channels. Voltage-gated calcium channels are activated resulting in a massive rise in intracellular calcium. Further exacerbating the problem is a large transient release of glutamate which itself

increases both Na^+ and Ca^{2+} influx through ionotropic glutamate receptors. Glutamate also binds to metabotropic receptors, which results in activation of the inositol phosphate pathway. This sets off a cascade of intracellular events, including further release of calcium from intracellular stores. It is now well accepted that this initial overload of intracellular calcium ultimately leads to the delayed cytotoxicity that is seen hours or days later.

Recently it has been reported that dopaminergic neurons exposed to a very short hypoxic challenge will hyperpolarize primarily through an opening of K_{ATP} channels (Guatteo et al., 1998). This stimulatory effect was suggested to be a direct result of the increased metabolic demand and the consequent drop in intracellular ATP levels. Furthermore Jovanovic et al. (1998b) recently reported that cells transfected with DNA encoding for Kir6.2/SUR1 (neuronal K_{ATP}) channel showed increased resistance to injury caused through hypoxia-reoxygenation. Therefore, the opening of K_{ATP} channels may serve a vital cytoprotective role during short periods of reduced oxygen in neuronal tissue. Thus, there is great therapeutic potential in developing compounds that not only will act to prevent this calcium influx prophylactically, but will aid in reestablishing the normal resting membrane potential in damaged tissue. Treatment with κ -PVIIA will act to open K_{ATP} channels, inducing membrane hyperpolarization and indirectly producing closure of the voltage-gated Ca^{2+} channels, thereby preventing or reducing deleterious effects of a massive calcium influx.

In accordance with the present invention, it has been found that intravenous (IV) injection of concentrations of κ -PVIIA, far higher than those required to produce maximal hyperpolarization in tracheal cultures *in vitro*, had no effect on blood pressure or heart rate in the anesthetized rat.

Our preliminary data indicates that kappa-PVIIA induces glibenclamide-sensitive currents in primary cultures of myocytes in a highly potent manner. Furthermore, incubation of primary myocyte cultures in the presence of κ -PVIIA confers protection against hypoxia-induced depolarization.

EXAMPLES

The present invention is described by reference to the following Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below were utilized.

EXAMPLE 1

Experimental Methods1. Cell culture protocol

Primary cultures of rat neonatal cortical cells, ventricular myocytes, tracheal smooth muscle cells and hippocampal cells were prepared. Cortical hemispheres were cleaned of meninges and the hippocampus removed and dissociated separately using 20 U/ml Papain. Cells were dissociated with constant mixing for 45 min at 37°C. Digestion was terminated with fraction V BSA (1.5 mg/ml) and Trypsin inhibitor (1.5 mg/ml) in 10 mls media (DMEM/F12 ± 10 % fetal Bovine serum ± B27 neuronal supplement; Life Technologies). Cells were gently triturated, to separate cells from surrounding connective tissue. Using a fluid-handling robot (Quadra 96, Tomtec) cells were settled onto Primaria-treated 96 well plates (Becton-Dickinson). Each well was loaded with approximately 25,000 cells. Plates were placed into a humidified 5% CO₂ incubator at 37°C and kept for at least five days before fluorescence screening. Ventricles were diced into 2mm square pieces and were digested in the presence of 20 U/ml Papain and trypsin/EDTA 1X (Life technologies). Smooth muscle cells on the surface of the trachea were cultured using the same digestive enzymes. Culturing techniques followed the method above.

2. Fluorimetry Assay

The saline solution used for the fluorimetric assay contained [in mM] 137 NaCl, 5 KCl, 10 HEPES, 25 Glucose, 3 CaCl₂, and 1 MgCl₂.

Di-8-ANEPPs: Voltage-sensitive dye. The effects of the compounds on membrane-potential were examined using the voltage-sensitive dye Di-8-ANEPPs. The Di-8-ANEPPs (2 µM) was dissolved in DMSO (final bath concentration 0.3%) and loaded into the cells in the presence of 10% pluronic acid. The plates were incubated for 40 min and then washed 4 times with the saline solution before starting the experiments. Di-8-ANEPPs crosses over the membrane in the presence of the pluronic acid creating a cytoplasmic pool of dye. Di-8-ANEPPs inserts into the plasma membrane where changes in potential result in molecular rearrangement. During hyperpolarization, the dye interchelates into the outer leaflet of the plasma membrane from the cytoplasmic reservoir of dye. Hyperpolarizations are represented as a positive shift and depolarizations as a negative shift in the fluorescence levels. ANEPs dyes show a fairly uniform 10% change in fluorescence intensity per 100mV change in membrane potential and as such, fluorescence changes can be correlated to changes in membrane potential.

PBFI:K⁺ sensitive dye. A lipid-soluble AM ester of the PBFI dye was used to examine the effect of the κ -PVIIA on intracellular potassium levels. The dye was loaded into the cytoplasm with 20 % pluronic acid where esterases cleave the dye from the ester effectively trapping the dye within the cell. Increases in intracellular potassium (K⁺) are reflected as a rise in fluorescence and decreases in K⁺ as a drop in fluorescence. Cells were pre-incubated in 5uM PBFI for three to four hours prior to screening. As with the Di-8-ANEPPs dye, the plates were rinsed four times with saline prior to beginning the experiments.

Fluo-3- Calcium-sensitive dye. To examine changes in intracellular calcium a lipid-soluble ester of the Fluo-3 dye (2uM in DMSO. Final bath concentration of DMSO 0.3%) is loaded into the cells in the presence of 20% pluronic acid. The plates are incubated for 35 minutes and washed four times with saline solution before beginning the experiments. Increases and decreases in the concentration of intracellular calcium are reflected as positive and negative changes in the percent fluorescence respectively.

Ethidium homodimer-1: cellular viability dye. The degree of cellular damage produced by a cytotoxic agent was measured using the dye Ethidium homodimer-1 (Molecular probes). This dye will not cross intact plasma membranes, but is able to readily enter damaged cells. Upon binding nucleic acids, the dye undergoes a fluorescent enhancement. Thus, the degree of cellular damage can be correlated to the amount of fluorescence. In preparation for the excitotoxicity assay, the cells were rinsed three times and pretreated with the kappa-PVIIA or an equal volume of saline. The cells were incubated for 15 minutes and glutamate (5-500uM) added to the appropriate lanes of the plate. The cells were incubated for a further 30 minutes, and washed thoroughly four times. The Ethidium Dye (4uM) was loaded into all the wells and a reading was taken immediately. Readings were then taken at hourly intervals.

3. Fluorimetry protocol

Fluorometric measurements are an averaging of cellular responses from approximately 25,000 cells per well of a 96 well plate. Cultures of cells from the cortex include at least pyramidal neurons, bipolar neurons, interneurons and astrocytes. Changes in membrane potential (Di-8-ANEPPs), cellular damage (Ethidium homodimer-1), intracellular K⁺ (PBFI) and Ca²⁺ (Fluo-3) were used as a measure of the response elicited with κ -PVIIA alone or with κ -PVIIA in the presence of specific receptor/ion channel agonists or antagonists. Concentration-responses were collected with the κ -PVIIA to determine the effective range. In order to minimize well-to-well variability, each

well acted as its own control by comparing the degree of fluorescence in pretreatment to that in post-treatment. This normalization process allows comparison of relative responses from plate to plate and culture to culture. Mixed-cell populations in each well were measured with the fluorimeter and individual cell signaling responses were averaged. Statistics, including mean and standard error of the mean, from eight wells allowed for comparison of significant differences between treatments. Results were expressed as percent change in fluorescence. An initial reading of a plate was taken in saline solution. Measurements using the Di-8-ANEPPs, Fluo-3 or PBFI dyes were made at time intervals of 15 seconds, two minutes, five minutes, 10 minutes, 20 minutes and 30 minutes in the presence of the compound. Readings with Ethidium homodimer-1 were made at hourly intervals.

4. Tracheal Smooth Muscle Preparation

Guinea pigs were sacrificed by cervical dislocation and the trachea excised and cleaned of connective tissue. Trachea were cut into four or five sections and opened by cutting through the ring of cartilage opposite the tracheal muscle. Each segment was mounted in an organ bath containing (mM) NaCl 118.2; KCl 4.7; $MgSO_4$ 1.2; KH_2PO_4 1.2; Glucose, 11.7; $CaCl_2$ 1.9 and $NaHCO_3$ 25.0. The bath was maintained at 37°C and gassed with 95% O_2 and 5% CO_2 . The preparation was maintained under 1 g of tension and equilibrated for 60 minutes before starting the experiment. Contractions were measured isometrically using a force-displacement transducer connected to a Grass polygraph. Following the 60 minutes equilibration period, the trachea were exposed to a submaximal concentration of histamine. This step was repeated until the contractile response to the spasmogen is consistent. The relaxant effects of increasing concentrations of kappa-PVIAA was determined in the absence and presence of the histamine.

5. Patch Clamp Recording

Whole-cell patch clamp recordings were made from cortical neurons on coverslips coated with Polyornithine/Poly-D-lysine (5 to 28 days in culture) and from myocytes on uncoated coverslips. Patch pipettes were pulled from thin-wall borosilicate glass and had resistances of 4M to 6M. Currents were recorded with an EPC 9 amplifier (HEKA) and controlled by software (Pulse, HEKA) run on a Macintosh power PC. Whole-cell currents were low-passed filtered at 10 kHz, digitized through a VR-10b digital data recorder to be stored on videotape at a sampling rate of 94 kHz. The intracellular pipette contained (in mM): 107 KCl, 33 KOH, 10 EGTA, 1 $MgCl_2$, 1 $CaCl_2$.

and 10 HEPES. The solution was brought to pH 7.2 with NaOH and 0.1-0.5 mM Na_2ATP and 0.1mM NaADP were added immediately before the experiment. The extracellular solution contained (in mM): 60 KCl, 80 NaCl, 1 MgCl_2 , 0.1 CaCl_2 and 10 HEPES. The pH of the external solution was brought to pH 7.4 with NaOH. The high concentration of potassium results in a calculated reversal potential for potassium of -20 mV. As a result, if the holding potential is more negative than -20mV, opening K channels will result in an inward flux of K ions and a downward deflection of the whole cell current. These solutions were chosen as the K_{ATP} channel has weak inward rectifying properties and as such, larger inward currents were anticipated. Experiments that are underway will address the effect of κ -PVIIA in solutions with low potassium levels.

6. Electrophysiology Solutions

Two extracellular solutions were used with different K^+ ion and Na^+ ion concentrations. Solution 1 contained 5 mM KCl and has a potassium equilibrium potential (E_K) of -84 mV, and solution 2 contained 60 mM and has a corresponding E_K of -20 mV. Extracellular solution 1 contained (in mM): 5 KCl, 135 NaCl, 1 MgCl_2 , 0.1 CaCl_2 and 10 HEPES. The pH of the external solution was corrected to pH 7.4 with NaOH. Extracellular solution 2 contained (in mM): 60 KCl, 80 NaCl, 1 MgCl_2 , 0.1 CaCl_2 and 10 HEPES. The pH of the external solution was corrected to pH 7.4 with NaOH. The intracellular pipette contained (in mM): 107 KCl, 33 KOH, 10 EGTA, 1 MgCl_2 , 1 CaCl_2 and 10 HEPES. The solution was brought to pH 7.2 with NaOH and 0.1-0.5 mM Na_2ATP , and 0.1mM NaADP was added immediately before the experiment.

7. Interpreting the Electrophysiology Results

In the presence of a low concentration of external K^+ ions (solution 1) and at holding potentials more depolarized than -84 mV, the opening of K^+ channels will result in an outward flux of K^+ ions. In the presence of a high concentration of K^+ (solution 2) the membrane potential would have to be more negative than -20 mV in order to see an outward movement of K ions. If the actual reversal potentials of the current evoked by κ -PVIIA in two different extracellular solutions are the same as the calculated values, it is highly likely that the κ -PVIIA-induced current is a result of the flux of K ions. The reversal potential of the current was calculated by holding the cell at the calculated E_K and running 500ms voltage ramps from -100mV to +80mV both in the presence and absence of increasing concentrations of κ -PVIIA. The average of four control ramps was subtracted from the average of four ramps evoked in the presence of κ -PVIIA. The resultant trace was the

actual current induced by the presence of the compound. This was fitted with a polynomial function and the reversal potential calculated.

8. Time-lapse Confocal Ca^{2+} Imaging

Cortical cell cultures were loaded with the fluorescent Ca^{2+} indicator Fluo3-AM (Molecular Probes, Eugene OR; 2mM final concentration with 0.1% Pluronic acid) 40 minutes prior to imaging experiments. Coverslips containing cells were mounted in a laminar flow perfusion chamber (Cornell-Bell design; Warner Instruments, Hamden, CT) and rinsed in saline (137 mM NaCl, 5 mM KCl, 3 mM CaCl_2 , 1 mM MgCl_2 , 10 mM HEPES, and 20 mM Sorbitol, pH 7.3) for at least five minutes to remove excess Fluo-3AM. Time-lapse images were collected on a Nikon PCM200 (Melville, NY) confocal scanning laser microscope equipped with a Zeiss Axiovert135 inverted microscope (Carl Zeiss, Inc., Thornwood, NY) and downloaded with no frame averaging every 1.8 seconds to an optical memory disk recorder (Panasonic TQ3031F, Secaucus NJ) (see methods further described in Kim et al., 1994). Image analysis were performed on a standardized 5×5 pixel area of cytoplasm in every astrocyte in the field to prevent bias in data analysis. Time course plots of intensity measurements (% change in fluorescence) were obtained using programs written by H. Sontheimer (Birmingham, AL) and plotted using Origin (MicroCal Northampton, MA). Routine analysis consisted of time course plots for up to 200 cells per field with at least five trials, thus yielding data analysis often from thousands of cells per experiment.

EXAMPLE 2

Exposure to κ -PVIIA Produces a Dose-Dependent Decrease in Intracellular K^+

κ -PVIIA was originally isolated from the purple cone snail (*Conus purpurascens*) and was found to block the *Drosophila* H4 *shaker* K^+ channel (Shon et al, 1998). In the same study no effects of the peptide were noted in oocytes expressing the mammalian *shaker*-like voltage-sensitive K^+ channels Kv1.1 and Kv1.3. The potential of the peptide to block other voltage-gated K^+ channels present in primary cultures of cortex was tested in this study. A 96-well fluorimetry assay was used to look for changes in potassium levels under depolarized conditions where voltage-gated potassium channels (Kv) would be activated. The cells were preloaded with the potassium indicator dye PBFI. If the compound acted to block Kv channels in a depolarized environment, there would be a resultant increase in intracellular K^+ . The results, however, suggested that at concentrations up to 100 nM, there was a reduction in the intracellular K^+ concentration in untreated resting preparations

(Figure 1), as well as those preparations depolarized with 10-100 μ M Aconitine. While the changes in fluorescence in the PBFI dye evoked with κ -PVIIA are small, it is important to stress that they are significant and repeatable.

EXAMPLE 3

Exposure to κ -PVIIA Produces Dose-Dependent Hyperpolarization

The fluorimetry experiments were repeated in the presence of the voltage-sensitive dye Di-8-ANEPPs, and the drop in intracellular K^+ levels was seen to be accompanied by a significant hyperpolarization of the preparation (represented by a positive shift in the fluorescence, Figures 2A-2B). κ -PVIIA is extremely potent in this assay, showing EC_{50} s of 8×10^{-16} M in cortex, 9×10^{-16} M in myocyte cultures and 9×10^{-18} M in primary cultures of tracheal myocytes.

EXAMPLE 4

The κ -PVIIA-Induced Hyperpolarization is Blocked by Exposure to K_{ATP} Antagonists

In order to determine the involvement of different K^+ channel subtypes in the κ -PVIIA-induced hyperpolarization, effects of five well-documented K^+ channel antagonists (4-aminopyridine (4-AP), Iberitoxin (IBTX), Apamin, Tolbutamide and Glibenclamide) were tested. In cortical preparations, applications of 4-AP, IBTX and Apamin were without any detectable effect on the hyperpolarization seen with 100 nM κ -PVIIA. However, both Tolbutamide (1-10 μ M) and Glibenclamide (10 nM), antagonists of the K_{ATP} channel, produced significant reductions in the κ -PVIIA induced hyperpolarization (Figure 3B). Glibenclamide also produced significant reductions in the κ -PVIIA-induced hyperpolarization in cultures of myocytes (Figure 3A).

EXAMPLE 5

κ -PVIIA Induces Tolbutamide or Glibenclamide-Sensitive Currents

The sensitivity of the response to K_{ATP} antagonists was confirmed using the whole-cell patch clamp technique. In these experiments, the extracellular potassium concentration was increased to 60 mM and the solutions were calculated such that the reversal potential for potassium (E_K) would be -20 mV. Thus, the opening of K^+ channels when the membrane potential is more negative than -20 mV will result in an influx of K^+ ions. In both primary cultures of cortex and cardiac myocytes, the superfusion of 100 nM κ -V11A induced an inward flux of positive ions that reversed close to -20 mV, indicating the involvement of K^+ ions. With a holding potential of -80 mV, the currents

evoked by κ -PVIIA were significantly larger in the myocyte preparation (87.7 ± 5.9 pA, $n=8$) compared to the cortical preparation (26.2 ± 6.2 pA, $n=4$). Even when the currents are corrected for cell capacitance, responses produced by the myocytes were greater than those seen in the cortical preparation (4.6 ± 0.4 pA/pf and 2.4 ± 0.7 pA/pf, respectively).

In both cases, the currents were sensitive either to the K_{ATP} antagonists tolbutamide (100 μ M) or glibenclamide (10 nM) (Figures 4A and B). The reversal potential of the κ -PVIIA evoked current was determined using a voltage ramp from -100 to +60 mV and fitting the results with a fourth-order polynomial fit (Figure 4C). The experimentally determined E_k (-23 mV) was close to the calculated E_k of -20 mV for these high potassium solutions, indicating the involvement of K^+ channels.

EXAMPLE 6

κ -PVIIA Produces a Slowly Developing Reduction in Intracellular Calcium

The effects of κ -PVIIA on intracellular calcium levels were determined using a 96-well fluorimetry assay plate and loading the cells with the Ca^{2+} indicator dye Fluo-3. In primary cultures of cortical neurons, κ -PVIIA produced a significant reduction in intracellular calcium. Little effect was noticeable with 1 nM κ -PVIIA at 15 seconds ($-2.15 \pm 0.95\%$, two trials) but over time, the drop in calcium concentration became more profound (30 min, $-8.8 \pm 3.9\%$).

EXAMPLE 7

κ -PVIIA Protects Against Hypoxia-Induced Depolarization

The depolarizing effects of N_2 -induced hypoxia have been monitored in cardiac ventricular myocytes using the voltage sensitive dye Di-8-ANEPPs in a 96 well fluorimetry assay plate. Solutions were depleted of oxygen by constant bubbling with N_2 gas and were compared to results with control untreated saline. Under these conditions, hypoxia produced significant depolarization of the preparation (reflected as a drop in fluorescence), and incubating the preparation with 10 nM κ -PVIIA prevented any hypoxia-induced changes in membrane potential (Figure 5).

EXAMPLE 8

κ -PVIIA Protects Against Glutamate-Induced Excitotoxicity

The protective effect of κ -PVIIA against glutamate-induced excitotoxicity was tested, using the 96-well fluorimetry assay and the Ethidium homodimer-1 dead cell dye. Five lanes of the 96-well plate were pre-exposed to 100 pM κ -PVIIA, and another five to control saline. Glutamate was

then applied for 30 minutes, at which time the entire plate was washed thoroughly to remove all κ -PVIIA and glutamate. Ethidium dye was loaded, an initial reading taken and the amount of delayed cytotoxicity monitored for six hours. Increases in fluorescence represent increased cell destruction. As can be seen from Figure 6, pre-incubating the cortical cells in κ -PVIIA resulted in very effective protection against the delayed (6 hrs) cytotoxic effects of 100uM glutamate. This protection was blocked by 100uM tolbutamide (K_{ATP} antagonist).

EXAMPLE 9

Cytotoxicity of κ -PVIIA

Incubation of primary cortical cultures with 200nM κ -PVIIA for 20 minutes induced no detectable protease activity (three trials). In comparison, a 20 minutes incubation with 5% Triton produced an ~14% increase in fluorescence, as detected by the Enzchek protease-sensitive dye.

EXAMPLE 10

Evaluation of κ -PVIIA as a Bronchodilator

The ability of κ -PVIIA to relax histamine-contracted, isolated Guinea pig tracheal segments is tested, using isometric tension recording. It is found that κ -PVIIA is able to relax histamine-contracted, isolated Guinea-pig tracheal segments. The response of κ -PVIIA is also tested in the presence of the K_{ATP} channel antagonists Tolbutamide or Glibenclamide. It is found that these antagonists reduce effects of κ -PVIIA, confirming involvement of the K_{ATP} channel in the response.

EXAMPLE 11

Evaluating Protective Ability of κ -PVIIA in *in vitro* Model of Hypoxia

A combination of the 96-well fluorimetric assay, electrophysiology, and confocal microscopy are used to assess the ability of κ -PVIIA to protect against the acute effects of transiently depleting oxygen in primary cultures. A multi-chamber saline reservoir has been constructed that allows the lower half of delivery plate to be filled with saline that is bubbled with N_2 . Individual chambers allow the effects of decreasing oxygen to be monitored in the presence and absence of different concentrations of the κ -PVIIA. An initial screen in primary cultures of ventricular myocytes, using the potentiometric dye Di-8-ANEPPs, shows a strong protective effect of the κ -PVIIA against hypoxia induced depolarization. Similar effects are seen in the cortex and trachea. When the

calcium-sensitive dye fluo-3 is used to observe changes in intracellular calcium levels induced by the hypoxic challenge, it is seen that κ -PVIIA is able to provide protection against hypoxia in all three tissue preparations. A similar result is obtained using the current-clamp mode of the whole cell patch clamp technique to monitor changes in membrane potential induced by hypoxia electrophysiology. This technique is very sensitive and allows the examination of the effect of κ -PVIIA on single tracheal, neuronal or myocyte cells.

EXAMPLE 12

Evaluating Protective Ability of κ -PVIIA in *in vitro* Model of Excitotoxicity

Preliminary fluorimetric experiments monitoring the degree of delayed cellular death produced following a challenge to a high concentration of glutamate have been carried out in primary cultures of cortex. The results indicate that the presence of the κ -PVIIA effectively reduces the degree of glutamate-induced excitotoxicity in a dose-dependant manner. Using the current-clamp mode of the whole-cell patch clamp technique, correlation of the fluorimetry results to actual changes in the membrane potential is examined. It is seen that the presence of the κ -PVIIA prevents the initial glutamate-induced depolarization, thereby conferring protection against the glutamate-induced calcium influx.

It will be appreciated that the methods and compositions of the instant invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will be apparent to the artisan that other embodiments exist and do not depart from the spirit of the invention. Thus, described embodiments are illustrative and should not be construed as restrictive.

LIST OF REFERENCES

- Ando, T. et al. (1997). *Clin. Exp. Allergy*. **27**:706-713.
- Barnay, G. et al. (2000). *J. Med. Chem.*
- Barnes, P.J. (1992). *Eur. Respir. J.* **5**:1126-1136.
- Bitan, G. et al. (1997). *J. Peptide Res.* **49**:421-426.
- Chen, J.W. et al. (1997) *Am. J. Cardiol.* **80**: 32-38.
- Ettinger, L.J. et al. (1978). *Cancer* **41**:1270-1273.
- Garcia, E. et al. (1999). *J. Gen. Physiol.* **114**:141-157.
- Goldschmidt, M. et al. (1996). *J. Clin. Pharmacol.* **36**:559-572.

- Guattee, E. et al. (1998). *J. Neurophysiol.* **79**:1239-1245.
- Hubry, V. et al. (1994). *Reactive Polymers* **22**:231-241.
- Jovanovic, A. et al. (1998a). *Circulation* **98**:1548-1555.
- Jovanovic, A. et al. (1998b). *Lab. Invest.* **78**:1101-1107.
- 5 Jung, Y.S. et al. (1998). *Japan J. Pharmacol.* **76**(1):65-73.
- Kim et al. (1994).
- Koike, A. et al. (1995). *Am J. Cardiol.* **76**:449-452.
- Kouchi, I. et al. (1998). *Am. J. Physiol.* **274**(4/2); H1106-H1112.
- Lauritzen I. et al. (1997). *J. Neurochem.* **69**(4); 1570-1579.
- 10 Lin, C. et al. (1998). *Pharmacology* **57**(6); 314-322.
- Luer, M.S. & Hatton, J. (1993). *Annals Pharmacotherapy* **27**:912-921.
- Morley, J. (1994). *Trends Pharmacol. Sci.* **15**(12); 463-468.
- Muller-Shweinitzer, E. and Fozard, J.R. (1997). *Br. J. Pharmacol.* **120**(7):1241-1248.
- Nagai, H. et al. (1991). *Japan J. Pharmacol.* **56**(1):13-21.
- 15 Neilson-Kudsk, J.E. (1996). *Dan. Med. Bull.* **43**(5): 429-447.
- Noma, A. (1983). *Nature* **304**:147-148.
- Ornstein, et al. (1993). *Biorganic Medicinal Chemistry Letters* **3**:43-48.
- Pell, T.J. et al. (1998). *Am. J. Physiol.* **275**(5 pt 2):H542-547.
- Rakel, R.E. (1997). Conn's current therapy.
- 20 Reshef A. et al. (1998). *Neurosci. Lett.* **250**(2):111-114.
- Sambrook, J. et al. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Shon, K.-J. et al. (1998). *J. Biol. Chem.* **273**:33-38.
- Terlau, H. et al. (1996). *Nature* **381**:148-151.
- 25 Terlau, H. et al. (1999). *J. Gen. Physiol.* **114**:125-140.
- Van de Steen, P. et al. (1998). *Critical Rev. in Biochem. and Mol. Biol.* **33**:151-208.
- Wind, T. et al. (1997). *Brain Res.* **751**(2):295-299.
- Yamabe H. et al. (1995). *Cardiovasc. Drugs Ther.* **9**(6):755-761.
- U.S. Patent No. 5,331,001.
- 30 U.S. Patent No. 5,534,615.
- U.S. Patent No. 5,364,769.

U.S. Patent No. 5,545,723.

U.S. Patent No. 5,550,050.

U.S. Patent No. 5,844,077.

U.S. Patent No. 5,672,682.

5 PCT Published Application WO 92/19195.

PCT Published Application WO 94/25503.

PCT Published Application WO 95/01203.

PCT Published Application WO 95/05452.

PCT Published Application WO 96/02286.

10 PCT Published Application WO 96/02646.

PCT Published Application WO 96/11698.

PCT Published Application WO 96/40871.

PCT Published Application WO 96/40959.

PCT Published Application WO 97/12635.

15 PCT Published Application WO 00/23092.

WHAT IS CLAIMED IS:

1. A method for treating disorders associated with radical depolarization of excitable membranes by activating a K_{ATP} channel which comprises administering to an individual in need thereof an effective amount of an active agent selected from the group consisting of:

(a) a compound of the following formula

Cys-Xaa₁-Ile-Xaa₂-Asn-Gln-Xaa₃-Cys-Xaa₄-Gln-Xaa₅-Leu-Asp-Asp-Cys-Cys-Ser-Xaa₁-Xaa₃-Cys-Asn-Xaa₁-Xaa₄-Asn-Xaa₃-Cys-Val (SEQ ID NO:1), wherein Xaa₁ and Xaa₃ are independently Arg, homoarginine, ornithine, Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys, any synthetic basic amino acid, His or halo-His; Xaa₂ is Pro or hydroxy-Pro (Hyp); Xaa₄ is Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; and Xaa₅ is His or halo-His,

(b) an analog of the compound of (a), said analog selected from the group consisting of:

κ-PVIIA[R18A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Ala-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:2);

κ-PVIIA[R22A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Ala-Phe-Asn-Lys-Cys-Val (SEQ ID NO:3);

κ-PVIIA[I3A]: Cys-Arg-Ala-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:4);

κ-PVIIA[K19A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Ala-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:5);

κ-PVIIA[R2A]: Cys-Ala-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:6);

κ-PVIIA[F9A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Ala-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:7);

κ-PVIIA[K25A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Ala-Cys-Val (SEQ ID NO:8);

κ-PVIIA[R2K]: Cys-Lys-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:9);

κ-PVIIA[K7A]: Cys-Arg-Ile-Hyp-Asn-Gln-Ala-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:10);
 κ-PVIIA[F9M]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Met-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:11);
 κ-PVIIA[F9Y]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Tyr-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:12);
 κ-PVIIA[R2Q]: Cys-Gln-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:13);
 κ-PVIIA[H11A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-Ala-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:14);
 κ-PVIIA[D14A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Ala-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:15);
 κ-PVIIA[Q6A]: Cys-Arg-Ile-Hyp-Asn-Ala-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:16);
 κ-PVIIA[N21A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Ala-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:17);
 κ-PVIIA[S17A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ala-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:18);
 κ-PVIIA[N24A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Ala-Lys-Cys-Val (SEQ ID NO:19);
 κ-PVIIA[L12A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Ala-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:20);
 κ-PVIIA[D13A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Ala-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:21);
 κ-PVIIA[Q10A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Ala-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:22);
 κ-PVIIA[V27A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Ala (SEQ ID NO:23);
 κ-PVIIA[O4A]: Cys-Arg-Ile-Ala-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:24); and
 κ-PVIIA[N5A]: Cys-Arg-Ile-Hyp-Ala-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-
 Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:25);

- (c) a derivative of (a) or (b); and
- (d) a physiologically acceptable salt thereof.

2. The method of claim 1, wherein Xaa₂ is hydroxy-Pro.
3. The method of claim 1, wherein Xaa₁ is Arg, Xaa₃ is Lys, Xaa₄ is Phe and Xaa₅ is His.
4. The method of claim 3, wherein Xaa₂ is hydroxy-Pro.
5. The method of claim 1, wherein said disorder is cardiac ischemia.
6. The method of claim 1, wherein said disorder is cerebral ischemia.
7. The method of claim 1, wherein said disorder is asthma.
8. The method of claim 1, wherein said disorder is ocular ischemia.
9. The method of claim 1, wherein the derivative is peptide of (a) or (b) in which the Arg residues may be substituted by Lys, ornithine, homoargine, nor-Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys or any synthetic basic amino acid; the Lys residues may be substituted by Arg, ornithine, homoargine, nor-Lys, or any synthetic basic amino acid; the Tyr residues may be substituted with any synthetic hydroxy containing amino acid; the Ser residues may be substituted with Thr or any synthetic hydroxylated amino acid; the Thr residues may be substituted with Ser or any synthetic hydroxylated amino acid; the Phe and Trp residues may be substituted with any synthetic aromatic amino acid; the Asn, Ser, Thr or Hyp residues may be glycosylated (contain an N-glycan or an O-glycan); the Cys residues may be in D or L configuration and may optionally be substituted with homocysteine (D or L); the Tyr residues may also be substituted with the 3-hydroxyl or 2-hydroxyl isomers (meta-Tyr or ortho-Tyr, respectively) and corresponding O-sulpho- and O-phospho-derivatives; the acidic amino acid residues may be substituted with any synthetic acidic amino acid, e.g., tetrazolyl derivatives of Gly and Ala; the aliphatic amino acids may be substituted by synthetic derivatives bearing non-natural aliphatic branched or linear side

chains C_nH_{2n+2} up to and including $n=8$; and pairs of Cys residues may be replaced pairwise with isoteric lactam or ester-thioether replacements, such as Ser/(Glu or Asp), Lys/(Glu or Asp) or Cys/Ala combinations.

$\mu^{(0)}$
 $\mu^{(1)}$
 $\mu^{(2)}$
 $\mu^{(3)}$
 $\mu^{(4)}$
 $\mu^{(5)}$
 $\mu^{(6)}$
 $\mu^{(7)}$
 $\mu^{(8)}$
 $\mu^{(9)}$
 $\mu^{(10)}$
 $\mu^{(11)}$
 $\mu^{(12)}$
 $\mu^{(13)}$
 $\mu^{(14)}$
 $\mu^{(15)}$
 $\mu^{(16)}$
 $\mu^{(17)}$
 $\mu^{(18)}$
 $\mu^{(19)}$
 $\mu^{(20)}$
 $\mu^{(21)}$
 $\mu^{(22)}$
 $\mu^{(23)}$
 $\mu^{(24)}$
 $\mu^{(25)}$
 $\mu^{(26)}$
 $\mu^{(27)}$
 $\mu^{(28)}$
 $\mu^{(29)}$
 $\mu^{(30)}$
 $\mu^{(31)}$
 $\mu^{(32)}$
 $\mu^{(33)}$
 $\mu^{(34)}$
 $\mu^{(35)}$
 $\mu^{(36)}$
 $\mu^{(37)}$
 $\mu^{(38)}$
 $\mu^{(39)}$
 $\mu^{(40)}$
 $\mu^{(41)}$
 $\mu^{(42)}$
 $\mu^{(43)}$
 $\mu^{(44)}$
 $\mu^{(45)}$
 $\mu^{(46)}$
 $\mu^{(47)}$
 $\mu^{(48)}$
 $\mu^{(49)}$
 $\mu^{(50)}$
 $\mu^{(51)}$
 $\mu^{(52)}$
 $\mu^{(53)}$
 $\mu^{(54)}$
 $\mu^{(55)}$
 $\mu^{(56)}$
 $\mu^{(57)}$
 $\mu^{(58)}$
 $\mu^{(59)}$
 $\mu^{(60)}$
 $\mu^{(61)}$
 $\mu^{(62)}$
 $\mu^{(63)}$
 $\mu^{(64)}$
 $\mu^{(65)}$
 $\mu^{(66)}$
 $\mu^{(67)}$
 $\mu^{(68)}$
 $\mu^{(69)}$
 $\mu^{(70)}$
 $\mu^{(71)}$
 $\mu^{(72)}$
 $\mu^{(73)}$
 $\mu^{(74)}$
 $\mu^{(75)}$
 $\mu^{(76)}$
 $\mu^{(77)}$
 $\mu^{(78)}$
 $\mu^{(79)}$
 $\mu^{(80)}$
 $\mu^{(81)}$
 $\mu^{(82)}$
 $\mu^{(83)}$
 $\mu^{(84)}$
 $\mu^{(85)}$
 $\mu^{(86)}$
 $\mu^{(87)}$
 $\mu^{(88)}$
 $\mu^{(89)}$
 $\mu^{(90)}$
 $\mu^{(91)}$
 $\mu^{(92)}$
 $\mu^{(93)}$
 $\mu^{(94)}$
 $\mu^{(95)}$
 $\mu^{(96)}$
 $\mu^{(97)}$
 $\mu^{(98)}$
 $\mu^{(99)}$

TITLE OF THE INVENTION

USES OF KAPPA-CONOTOXIN PVIIA

ABSTRACT OF THE DISCLOSURE

The invention relates to uses of kappa-conotoxin PVIIA (κ -PVIIA), analogs and derivatives
 5 for activating ATP-sensitive K^+ channels. The activation of ATP-sensitive K^+ channels is useful
 for opening K_{ATP} channels which can be used to treat a wide range of disease and injury states,
 including cerebral and cardiac ischemia and asthma.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
219

Effect of K-PVIIA on potassium levels in primary cultures of myocytes.
15 min after K-PVIIA application (PBF dye)




FIG. 1

FIG. 2A Myocytes




FIG. 2B Cortex




FIG. 3A

Myocytes

FIG. 3B

Cortex





FIG. 4A)




FIG. 4B) Repeated applications of 100nM KPV11A





FIG. 4C)



N₂-induced hypoxia




FIG. 5




FIG. 6

SEQUENCE LISTING

<110> Cornell-Bell, Ann H.
 Pemberton, Karen E.
 Temple Jr., Davis L.
 Layer, Richard T.
 McCabe, R. Tyler
 Jones, Robert M.
 Cognetix, Inc.

<120> Uses of Kappa-Conotoxin PVIIA

<130> Kappa-PVIIA

<140>
 <141>

<150> US 60/219,438
 <151> 2000-07-20

<150> US 60/155,135
 <151> 1999-09-22

<160> 25

<170> PatentIn Ver. 2.0

<210> 1
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa at residue 2, 7, 18, 19, 22 and 25 may be Arg,
 homoarginine, ornithine, Lys, N-methyl-Lys,
 N,N-dimethyl-Lys, N,N,N-trimethyl-Lys, any
 synthetic basic amino acid, His or halo-His; Xaa
 at residue

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> 4 may be Pro or Hyp; Xaa at residue 9 and 23 may
 be Phe,Tyr, meta-Tyr, ortho-Tyr, nor-Tyr,
 mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr,
 O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp,

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> halo-Trp (D or L) or any synthetic aromatic amino
 acid; Xaa at residue 11 is His or halo-His

<400> 1
 Cys Xaa Ile Xaa Asn Gln Xaa Cys Xaa Gln Xaa Leu Asp Asp Cys Cys
 1 5 10 15

Ser Xaa Xaa Cys Asn Xaa Xaa Asn Xaa Cys Val
 20 25

<210> 2

<211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 2
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Ala Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 3
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 3
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Ala Phe Asn Lys Cys Val
 20 25

<210> 4
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 4
 Cys Arg Ala Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 5
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 5

Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
1 5 10 15

Ser Arg Ala Cys Asn Arg Phe Asn Lys Cys Val
20 25

<210> 6
<211> 27
<212> PRT
<213> Conus purpurascens

<220>
<221> PEPTIDE
<222> (1)..(27)
<223> Xaa is Hyp

<400> 6
Cys Ala Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
20 25

<210> 7
<211> 27
<212> PRT
<213> Conus purpurascens

<220>
<221> PEPTIDE
<222> (1)..(27)

<400> 7
Cys Arg Ile Xaa Asn Gln Lys Cys Ala Gln His Leu Asp Asp Cys Cys
1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
20 25

<210> 8
<211> 27
<212> PRT
<213> Conus purpurascens

<220>
<221> PEPTIDE
<222> (1)..(27)
<223> Xaa is Hyp

<400> 8
Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Ala Cys Val
20 25

<210> 9
<211> 27
<212> PRT
<213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 9
 Cys Lys Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 10
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 10
 Cys Arg Ile Xaa Asn Gln Ala Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 11
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 11
 Cys Arg Ile Xaa Asn Gln Lys Cys Met Gln His Leu Asp Asp Cys Cys
 1 5 10 15
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 12
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 12
 Cys Arg Ile Xaa Asn Gln Lys Cys Tyr Gln His Leu Asp Asp Cys Cys
 1 5 10 15
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 13
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 13
 Cys Gln Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 14
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 14
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln Ala Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 15
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)

<400> 15
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Ala Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 16
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 16
 Cys Arg Ile Xaa Asn Ala Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 17
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 17
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Ala Arg Phe Asn Lys Cys Val
 20 25

<210> 18
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 18
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ala Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 19
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 19
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Ala Lys Cys Val
 20 25

<210> 20
 <211> 27

<212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)

<400> 20
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Ala Asp Asp Cys Cys
 1 5 10 15
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 21
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 21
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Ala Asp Cys Cys
 1 5 10 15
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 22
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 22
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Ala His Leu Asp Asp Cys Cys
 1 5 10 15
 Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 23
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 23
 Cys Arg Ile Xaa Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Ala
 20 25

<210> 24
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<400> 24
 Cys Arg Ile Ala Asn Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25

<210> 25
 <211> 27
 <212> PRT
 <213> Conus purpurascens

<220>
 <221> PEPTIDE
 <222> (1)..(27)
 <223> Xaa is Hyp

<400> 25
 Cys Arg Ile Xaa Ala Gln Lys Cys Phe Gln His Leu Asp Asp Cys Cys
 1 5 10 15

Ser Arg Lys Cys Asn Arg Phe Asn Lys Cys Val
 20 25